



Forest / Forêt

Association CNVC00007

***Tsuga heterophylla* - *Thuja plicata* (*Pseudotsuga menziesii* - *Abies amabilis*) /
Gymnocarpium dryopteris - *Clintonia uniflora***

Western Hemlock - Western Red Cedar (Douglas-fir - Pacific Silver Fir) / Common Oak Fern - Singleflower
Clintonia

Pruche de l'Ouest - Thuya géant (Douglas de Menzies - Sapin gracieux) / Gymnocarpe du chêne - Clintonie
uniflore

Subassociations: none

CNVC Alliance: not yet determined

CNVC Group: not yet determined

Type Description

Concept: This mature to old-growth, coniferous forest association is endemic to the coast / interior climatic transition of the middle to leeward eastern valley systems of the Coast Mountains and the Fraser River valley of British Columbia. It occurs on various topographic positions with mainly steep slope gradients. Mesic, nutrient-rich soils are normal for CNVC00007; soil drainage is variable, ranging from rapid to moderately well. Western hemlock (*Tsuga heterophylla*) and western red cedar (*Thuja plicata*) co-dominate the canopy, often with Douglas-fir (*Pseudotsuga menziesii*) and Pacific silver fir (*Abies amabilis*). The shrub layer is relatively sparse, comprising mostly regenerating canopy trees with various blueberries: red huckleberry (*Vaccinium parvifolium*), Alaskan blueberry (*V. alaskaense*), and oval-leaved blueberry (*V. ovalifolium*). The diverse herb layer is dominated by three-leaved foamflower (*Tiarella trifoliata*), common oak fern (*Gymnocarpium dryopteris*) and singleflower clintonia (*Clintonia uniflora*). Lanky moss (*Rhytidiadelphus loreus*) and stairstep moss (*Hylocomium splendens*) are the main moss-layer species.

Vegetation: In these mature to old coniferous forests, *Tsuga heterophylla* and *Thuja plicata* dominate the canopy coverage, followed by *Pseudotsuga menziesii*, and often with *Abies amabilis*. Dry summer weather has contributed to a history of extensive wildfires, which in turn, has favoured the widespread distribution of *Pseudotsuga menziesii*. The shrub layer is relatively sparse, with regenerating *Tsuga heterophylla*, *Thuja plicata*, and *Abies amabilis* typically most abundant. *Vaccinium parvifolium*, *V. alaskaense*, *V. ovalifolium* and *Oplopanax horridus* are also commonly present, but with low cover. Although the herb layer often has only moderate cover, it can be diverse. Common herb layer species include *Tiarella trifoliata* (see the Comments section), *Gymnocarpium dryopteris*, *Streptopus lanceolatus*, *Polystichum munitum*, *Dryopteris expansa*, *Clintonia uniflora* and *Cornus canadensis*. *Rhytidiadelphus loreus*, *Hylocomium splendens* and *Plagiomnium* / *Rhizomnium* spp. are the main mosses in the moderately well-developed bryophyte layer.



***Tsuga heterophylla* - *Thuja plicata* (*Pseudotsuga menziesii* - *Abies amabilis*) /
Gymnocarpium dryopteris - *Clintonia uniflora* CNVC00007**

Type Description (cont'd)

Environment: CNVC00007 is found in a coastal maritime / interior continental transitional climate. In southern portions of the range, it is found at elevations from approximately 650 to 1200 mASL in the westerly drainages experiencing some maritime influence; in drainages farther to the east which experience less maritime influence, elevations range between approximately 900 and 1350 m. In the more northerly or central areas of distribution, much lower elevations are observed - from sea level to 700 m. The rainshadow effects of mountain ranges to the west contribute to summer-dry conditions which can lead to high wildfire hazard. Topositions for this association vary from upper to lower slopes. Soils are often developed on fluvial and colluvial materials, with textures ranging from loamy to sandy. Rapidly to moderately well-drained, mesic, nutrient-rich soils are normal for these sites. Forest floor materials consist largely of moderately thick (10-12 cm) mors.

Dynamics: An intensive logging history means that few of the once extensive old-forest successional stages remain. Under natural conditions, when wind exposure is indirect, stand replacement may be gradual through the mortality of individual or small numbers of canopy trees. Unless windthrow and fire have been pervasive, the age composition of these potentially old-forest stages tends to be uneven. Stand-replacing fires can occur in dry summer weather. *Armillaria* root disease (*Armillaria ostoyae*) is a high risk for *Abies amabilis* and a medium risk for *Pseudotsuga menziesii* and *Tsuga heterophylla*. *Annosus* root disease (*Heterobasidion annosum*) is a medium risk for *Abies amabilis*.

Range: CNVC00007 is found in British Columbia in higher elevation drainages of the Fraser River, both east and north from Chilliwack in the lower Fraser Valley, and in drainages on the eastern side of the Coast Mountains from valleys adjacent to the northern half of Harrison Lake northwestward as far as the lower Kimsquit River valley near the headwaters of Dean Channel, approximately 75 km north of Bella Coola. It is a Canadian endemic association.

Conservation Status (NatureServe)

Global Conservation Rank: GNR

National Conservation Rank: not yet determined

Subnational Conservation Rank: no applicable rank



Forest / Forêt

Association CNVC00007

***Tsuga heterophylla* - *Thuja plicata* (*Pseudotsuga menziesii* - *Abies amabilis*) /
Gymnocarpium dryopteris - *Clintonia uniflora***

Western Hemlock - Western Red Cedar (Douglas-fir - Pacific Silver Fir) / Common Oak Fern - Singleflower
 Clintonia

Pruche de l'Ouest - Thuya géant (Douglas de Menzies - Sapin gracieux) / Gymnocarpe du chêne - Clintonie
 uniflore

Distribution

Countries: Canada

Provinces / Territories / States: British Columbia

Ecozones and Ecoregions of Canada: Pacific Maritime: Coastal Gap, Pacific Ranges


Rowe's Forest Regions and Sections: Coast: Southern Pacific Coast, Northern Pacific
 Coast

Commission for Environmental Cooperation Ecological Regions of North America:
 Marine West Coast Forests

The Nature Conservancy (USA) and Nature Conservancy of Canada Ecoregions:
 North Cascades, S.E. Alaska - B.C. Coastal Forest and Mountains

Biogeoclimatic Ecosystem Classification of British Columbia (zones and subzones):
 CWH ms

Ecoregion Classification System of British Columbia (ecosections): Eastern Pacific
 Ranges, Kimsquit Mountains, Northern Pacific Ranges



Corresponding Types and Associations

CNVC00007	British Columbia	CWH ms 1 /04	<i>Abies amabilis</i> - <i>Streptopus amplexifolius</i> - <i>Streptopus lanceolatus</i> (in part)
		CWH ms 2 /04	<i>Abies amabilis</i> - <i>Streptopus amplexifolius</i> - <i>Streptopus lanceolatus</i> (in part)



Forest / Forêt

Association CNVC00007

***Tsuga heterophylla* - *Thuja plicata* (*Pseudotsuga menziesii* - *Abies amabilis*) /
Gymnocarpium dryopteris - *Clintonia uniflora***

Western Hemlock - Western Red Cedar (Douglas-fir - Pacific Silver Fir) / Common Oak Fern - Singleflower
 Clintonia

Pruche de l'Ouest - Thuya géant (Douglas de Menzies - Sapin gracieux) / Gymnocarpe du chêne - Clintonie
 uniflore

Vegetation Summary*

Association

CNVC00007

22 plots

Species Name [†]	%	%
	Cover	Presence

Overstory Trees

<i>Tsuga heterophylla</i>	35	95
<i>Thuja plicata</i>	30	82
<i>Pseudotsuga menziesii</i>	32	59
<i>Abies amabilis</i>	26	41
<i>Picea sitchensis</i>	10	27
Tree Stratum Cover (P₁₀ P₂₅ Mean P₇₅ P₉₀)[‡]	(65 70 75 80 85)	

Understory Woody Shrubs and Regenerating Trees

<i>Oplopanax horridus</i>	2	91
<i>Tsuga heterophylla</i>	12	77
<i>Vaccinium parvifolium</i>	2	77
<i>Vaccinium ovalifolium</i>	3	68
<i>Vaccinium alaskaense</i>	7	59
<i>Menziesia ferruginea</i>	1	55
<i>Thuja plicata</i>	7	50
<i>Abies amabilis</i>	11	45
<i>Paxistima myrsinites</i>	1	36
<i>Ribes lacustre</i>	2	32
<i>Rubus parviflorus</i>	2	32
<i>Acer circinatum</i>	3	27
<i>Rubus spectabilis</i>	3	27
<i>Picea sitchensis</i>	5	23
<i>Viburnum edule</i>	1	23
<i>Sambucus racemosa</i>	1	23
Shrub Stratum Cover (P₁₀ P₂₅ Mean P₇₅ P₉₀)[‡]	(5 10 25 42 49)	

Understory Herbs and Dwarf Shrubs

<i>Gymnocarpium dryopteris</i>	14	91
<i>Clintonia uniflora</i>	9	82
<i>Dryopteris expansa</i>	3	82
<i>Cornus canadensis</i>	6	73
<i>Streptopus lanceolatus</i>	4	73
<i>Goodyera oblongifolia</i>	1	73
<i>Polystichum munitum</i>	4	68
<i>Tiarella trifoliata</i> var. <i>unifoliata</i>	4	68



***Tsuga heterophylla* - *Thuja plicata* (*Pseudotsuga menziesii* - *Abies amabilis*) /
Gymnocarpium dryopteris - *Clintonia uniflora* CNVC00007**

Vegetation Summary (cont'd)*

Species Name [†]	Association CNVC00007	
	% Cover	% Presence
<i>Athyrium filix-femina</i>	2	68
<i>Orthilia secunda</i>	1	59
<i>Tiarella trifoliata</i>	6	55
<i>Rubus pedatus</i>	2	45
<i>Maianthemum racemosum</i>	1	45
<i>Maianthemum stellatum</i>	3	41
<i>Chimaphila menziesii</i>	1	41
<i>Prosartes hookeri</i>	2	36
<i>Actaea rubra</i>	1	36
<i>Galium triflorum</i>	0	36
<i>Asarum caudatum</i>	2	32
<i>Streptopus amplexifolius</i>	1	32
<i>Linnaea borealis</i>	1	27
<i>Moneses uniflora</i>	1	27
<i>Adiantum aleuticum</i>	2	23
<i>Chimaphila umbellata</i>	1	23
<i>Pyrola asarifolia</i>	1	23
<i>Corallorhiza maculata</i>	1	23
<i>Viola orbiculata</i>	0	23
Herb Stratum Cover (P₁₀ P₂₅ Mean P₇₅ P₉₀)[‡]	(15 23 45 64 80)	
Bryophytes and Lichens		
<i>Rhytidiadelphus loreus</i>	12	82
<i>Hylocomium splendens</i>	23	77
<i>Plagiomnium insignne</i>	5	50
<i>Eurhynchium oregonum</i>	2	50
<i>Plagiochila asplenioides</i>	4	45
<i>Rhytidiopsis robusta</i>	9	41
<i>Plagiothecium undulatum</i>	1	32
<i>Rhizomnium glabrescens</i>	4	27
<i>Rhytidiadelphus triquetrus</i>	2	27
Bryo-Lichen Stratum Cover		
(P₁₀ P₂₅ Mean P₇₅ P₉₀)[‡]	(15 20 44 64 89)	

* species present in > 20% of sample plots are listed

[†] see **Botanical Nomenclature** link at <http://cnvc-cnvc.ca> for botanical sources, synonyms and common names

[‡] P_x = Xth percentile (e.g., P₁₀ = 10th percentile)



Forest / Forêt

Association CNVC00007

Tsuga heterophylla - *Thuja plicata* (*Pseudotsuga menziesii* - *Abies amabilis*) /
Gymnocarpium dryopteris - *Clintonia uniflora*

Western Hemlock - Western Red Cedar (Douglas-fir - Pacific Silver Fir) / Common Oak Fern - Singleflower
Clintonia

Pruche de l'Ouest - Thuya géant (Douglas de Menzies - Sapin gracieux) / Gymnocarpe du chêne - Clintonie
uniflore

Site / Soil Characteristics

Association
CNVC00007
22 plots

Elevation Range (min–mean–max meters)

10–388–1228

Slope Gradient (% frequency)

very steep (5)
steep (41)
moderately steep (23)
gentle (18)
level (14)

Aspect (% frequency)

north (27)
east (23)
south (5)
west (32)
level (14)

Meso Toposition (% frequency)

crest / upper (18)
mid (36)
lower / toe (9)
level (18)
missing data (18)

Moisture Regime (% frequency)

mesic (68)
moist (32)

Nutrient Regime (% frequency)

poor (5)
medium (27)
rich (64)
saline (5)



Tsuga heterophylla - *Thuja plicata* (*Pseudotsuga menziesii* - *Abies amabilis*) /
Gymnocarpium dryopteris - *Clintonia uniflora* CNVC00007

Site / Soil Characteristics (cont'd)

Association
CNVC00007

Soil Parent Material (% frequency)

colluvium (27)
fluvial (23)
glaciomarine (5)
missing data (45)

Soil Rooting Zone Substrate (% frequency)

non-soil (27)
sandy (36)
coarse loamy (18)
silty (5)
organic (5)
missing data (9)

Root Restricting Depth (% frequency)

0 – 20 cm (9)
21 – 99 cm (9)
missing data (82)

Humus Form (% frequency)

mor (50)
moder (36)
missing data (14)



Forest / Forêt

Association CNVC00007

***Tsuga heterophylla* - *Thuja plicata* (*Pseudotsuga menziesii* - *Abies amabilis*) /
Gymnocarpium dryopteris - *Clintonia uniflora***

Western Hemlock - Western Red Cedar (Douglas-fir - Pacific Silver Fir) / Common Oak Fern - Singleflower
Clintonia

Pruche de l'Ouest - Thuya géant (Douglas de Menzies - Sapin gracieux) / Gymnocarpe du chêne - Clintonie
uniflore

Additional Characteristics

Species of High Conservation Concern:

Non-native Species:

Management Issues:

Type Statistics

Internal Similarity:

Confidence: high

Strength:

Related Concepts

Similar CNVC Associations: CNVC00028 *Tsuga heterophylla* - *Abies amabilis* / *Oplopanax horridus* / *Gymnocarpium dryopteris* Forest

Related United States National Vegetation Classification Associations: C EGL002837 *Tsuga heterophylla* - *Thuja plicata* (*Abies amabilis*) /
Cornus canadensis Forest

Relationships with Other Classifications:

Comments

On rich sites in this climate, the wetter CNVC00028 [*Tsuga heterophylla* - *Abies amabilis* / *Oplopanax horridus* / *Gymnocarpium dryopteris*] typically has *Rubus spectabilis* present as well as an abundance of *Oplopanax horridus*; when *Oplopanax horridus* is present in communities of CNVC00007, it is with low cover.

Tiarella trifoliata (three-leaved foamflower) may include *T. trifoliata* var. *laciniata* (cut-leaved foamflower), *T. trifoliata* var. *trifoliata* (three-leaved foamflower) and/or *T. trifoliata* var. *unifoliata* (one-leaved foamflower).

Source Information

Number of source plots for CNVC00007: 22

Information Sources: British Columbia Ministry of Forests and Range, Research Branch BECMaster database, October 2007 (22 plots)

Concept Authors: D. Meidinger, C. Chappell, C. Cadrin, G. Kittel, C. McCain, K. Boggs, J. Kagan, G. Cushon, A. Banner and T. DeMeo

Description Authors: D. Meidinger, A. Inselberg, C. Cadrin and K. Baldwin

Date of Concept: November, 2005

Date of Description: March, 2011



***Tsuga heterophylla* - *Thuja plicata* (*Pseudotsuga menziesii* - *Abies amabilis*) /
Gymnocarpium dryopteris - *Clintonia uniflora* CNVC00007**

Source Information (cont'd)

Classification References:

British Columbia Ministry of Forests and Range, Research Branch. 2007. Vegetation classification hierarchy: BECMaster database (October 2007). B.C. Min. For., Victoria, BC.

Meidinger, D.; Chappell, C.; Cadrin, C.; Kittel, G.; McCain, C.; Boggs, K.; Kagan, J.; Cushon, G.; Banner, A.; DeMeo, T. 2005. International vegetation classification of the Pacific Northwest: International correlation of temperate coastal forest plant associations of Oregon, Washington, British Columbia and Alaska. Contributors: B.C. Ministry of Forests, USDA Forest Service, B.C. Conservation Data Centre, Alaska Natural Heritage Program, Washington Natural Heritage Program, Oregon Natural Heritage Information Center.

Characterization References:

British Columbia Conservation Data Centre. 2007. B.C. Species and Ecosystems Explorer. B.C. Min. of Environ. Victoria, BC. Available: <http://www.env.gov.bc.ca/cdc/access.html> (accessed August 12, 2008).

British Columbia Ministry of Forests and Range, Research Branch. 2007. BECMaster database (October 2007). B.C. Min. For., Victoria, BC.

Green, R.N.; Klinka, K. 1994. A field guide to site identification and interpretation for the Vancouver Forest Region. B.C. Min. For., Res. Branch, Victoria, BC. Land Manage. Handb. No. 28. 285 p.

NatureServe. 2007. NatureServe Explorer: An online encyclopedia of life [web application]. Version 6.2. NatureServe. Arlington, VA, USA. Available: <http://www.natureserve.org/explorer> (accessed November 26, 2007).

The information contained in this factsheet is based on data and expert knowledge that is current to the date of description. As new information becomes available, the factsheet will be updated.

Suggested Citation: Meidinger, D.; Inselberg, A.; Cadrin, C.; Baldwin, K. *Tsuga heterophylla* - *Thuja plicata* (*Pseudotsuga menziesii* - *Abies amabilis*) / *Gymnocarpium dryopteris* - *Clintonia uniflora* [online]. Sault Ste. Marie, Ontario, Canada: Canadian National Vegetation Classification. March, 2011; generated Mar-28-2011; cited ENTER DATE ACCESSED. 9 p. Canadian National Vegetation Classification Association: CNVC00007. Available from <http://cnvc-cnvc.ca>. System Requirements: Adobe Acrobat Reader v. 7.0 or higher. ISSN 1916-3266.